

- Barhoumi W, Qualls WA, Archer R, Fuller DO, Chelbi I, Cherni S, Derbali M, Arheart KL, Elyes Zhioua E, Beier JC (2015) Irrigation in the arid regions of Tunisia impacts the abundance and apparent density of sand fly vectors of *Leishmania infantum*. *Acta Tropica* 141:73-78.
- Jiang, J, Fuller DO, Teh, SY, Koh, HL, DeAngelis, DL, Sternberg LSL (2015) Bistability of mangrove forests and competition with fresh water plants. *Agricultural and Forest Meteorology* (in press).
- Samson DM, Archer R, Qualls, WA, Alimi TO, Arheart KK, Fuller DO (2015) Environmental Assessment of Mosquito Ecology in Northern Haiti. *Journal of Vector Ecology* (in press).
- WM Monteiro, Val, FA, Siqueira, AM, Franca, GP, Sampaio VS, Melo GC, de Almeida ACG, Brito M, Peixoto HM, Fuller DO, Bassat Q, Romero GAS, Oliveira MRF, Lacerda MVG (2014) G6PD deficiency in Latin America: prevalence, variants and implications for malaria elimination. *Memórias do Instituto Oswaldo Cruz* 1-16:2014.
- Fuller DO, Troyo A, Alimi T, Beier JC (2014) Participatory Risk Mapping of Malaria Vector Exposure in Northern South America using Environmental and Population Data. *Applied Geography* 48:1-7.
- Conley AK, Fuller DO, Hadad N, Hassan AN, Gad AM, Beier JC (2014) Modeling the distribution of the West Nile and Rift Valley Fever vector *Culex pipiens* in arid and semi-arid regions of the Middle East and North Africa. *Parasites & Vectors* 7:289.
- Chang AY, Fuller DO, Carrasquillo O, Beier JC (2014) Social justice, climate change, and dengue. *Health and Human Rights* 16:93-104.
- Fuller DO, Wang Y (2014) Recent trends in satellite vegetation index observations indicate decreasing vegetation biomass in the southeastern saline Everglades wetlands. *Wetlands* 34:67-77.
- Barr, J.G., Engel, V., Fuentes, J.D., Fuller, D.O., and Kwon, H. (2013). Modeling light use efficiency in a subtropical mangrove forest equipped with CO2 eddy covariance. *Biogeosciences* 10:2145-2158.
- Fuller, D.O., Miettinen, J., and Meijaard E. (2013). Deforestation, degradation, and forest regrowth in Indonesia's protected areas from 2000-2010. *Indonesian Journal of Conservation* 2:1-13.

Fuller, D.O., Parenti, M.S., Hassan A.N., and Beier, J.C. (2012). Linking land cover and species distribution models to project potential ranges of malaria vectors: an example using *Anopheles arabiensis* in Sudan and Upper Egypt. *Malaria Journal* 11:264, doi:10.1186/1475-2875-11-264.

Fuller, D.O., Ahumada, M., Quiñones, M.L., Herrera, S., and Beier, J.C. (2012). Near-present and future distribution of *Anopheles albimanus* in Mesoamerica and the Caribbean Basin modeled with climate and topographic data. *International Journal of Health Geographics* 11:13.

Fuller, D.O., Parenti, M.S., Gad, A., and Beier, J.C. (2012). Land cover in Upper Egypt assessed using regional and global land cover products derived from MODIS imagery. *Remote Sensing Letters* 3: 171-180, doi: 10.1080/01431161.2011.551847.

Herrera S, Quiñones ML, Quintero JP, Corredor V, Fuller DO, Mateus JC, Calzada JE, Gutierrez, JB, Llanos A, Soto E, Menendez C, Wuk Y, Alonso P., Carrasquilla G, Galinski M, Beier JC Arevalo-Herrera M. (2012). Prospects for malaria elimination in non-Amazonian regions of Latin America. *Acta Tropica* 121: 315-323.

LeComber, S., Rossmo, K., Hassan, A., Fuller, D.O., Beier J.C. (2011). Geographic profiling as a novel spatial tool for targeting infectious disease control. *International Journal of Health Geographics* 10:35 doi:10.1186/1476-072X-10-35.

Fuller, D.O., Hardiono, M., and Meijaard, E. (2011). Deforestation projections for carbon-rich peat swamp forests of Central Kalimantan, Indonesia. *Environmental Management* 48: 436-447.

Wang, X., Fuller, D.O., Sternberg, L., and Miralles, F. (2011). Foliar nutrient accumulation in subtropical tree islands: A new chemohydrodynamic link between satellite vegetation indices and foliar $\delta^{15}\text{N}$ values. *Remote Sensing of Environment* 115: 923-930, doi:10.1016/j.rse.2010.11.019.